

CARPA Standard Treatment Manual 5th edition

Erratum

- Page 245: Medicines for chronic kidney disease
Step 3; Dot point 1; Indented dot point 1
Perindopril-indapamine should read **Perindopril-indapamide**
- Page 276: Syphilis and donovanosis
Dot Point 1; Indented dot point 1
Benzathine penicillin 1.8mg (2.4 million units) should read **Benzathine penicillin** 1.8g (2.4 million units)
- Page 358: Do – for infected testes

For all men with discharge from penis AND any Indigenous man aged under 45 years or non-Indigenous man under 35 years with no discharge

- Treat as STI – give
 - **Ceftriaxone** IM 250mg single dose
 - **AND Azithromycin** oral 1g single dose
 - **AND Doxycycline** (over 8 years) 200mg daily for 14 days

For any Indigenous man 45 years or over or non-Indigenous man 35 years or over with no discharge from penis

- Treat as UTI – give
 - **Cephalexin** 1g twice a day (bd) for 14 days
 - **OR Amoxicillin-clavulanate** 875/125mg twice a day (bd) for 14 days

Should read

For all men with discharge from penis AND any man under 45 years with no discharge

- Treat as sexually acquired – give
 - **Ceftriaxone** IM 500mg single dose
 - **AND Azithromycin** oral 1g single dose
 - **AND Doxycycline** (over 8 years) 200mg daily for 14 days

For any man 45 years or over with no discharge from penis

- Treat as UTI related – give
 - **Trimethoprim** daily for 14 days (*doses p404*)
 - **OR Nitrofurantoin** (not if chronic kidney disease stage 3, 4, 5) for 14 days (*doses p401*)
 - **OR Amoxicillin-clavulanate** for 14 days (*doses p394*)

- Page 404: Antibiotics Table
Trimethoprin-sulphamethoxazole (co-trimoxazole); 12 years and over
1600mg (1 tablet) should read 160mg (1 tablet)